

# The Effect of Fish Oil, Vitamin D and Protein on URTI Incidence in Young Active People

## Authors

M. Da Boit, B. M. Gabriel, P. Gray, S. R. Gray

## Affiliation

Institute of Medical Sciences, University of Aberdeen, Aberdeen, United Kingdom

## Key words

- immune
- exercise
- nutrition

## Abstract

Upper respiratory tract infections (URTI) are a frequent illness among athletes. We investigated the effect of a multi-nutrient supplement (vitamin D, fish oil and protein) on the occurrence of URTI in young active people. 42 young recreational athletes were randomly assigned to receive either supplementation (550 mg DHA, 550 mg EPA, 10 µg vitamin D3 and 8 g whey protein) or placebo for 16 weeks. Unstimulated saliva samples were collected by passive drool. Samples were analysed for IgA (sIgA) concentration and the secretion rate extrapolated by multiplying concentration by saliva flow rate. Physical activ-

ity levels and URTI incidence were monitored by questionnaire. Training status was not different between the 2 groups. There were no differences in the incidence, severity and duration of URTI. However the number of symptom days was lower in the supplemented compared to the control group ( $1.72 \pm 1.67$  vs.  $2.79 \pm 1.76$ ;  $P < 0.05$ ). sIgA concentration and secretion rate did not differ between groups. This study demonstrates that 16 weeks of supplementation with fish oil, vitamin D and protein did not modify the incidence, severity and duration of URTI, although the total number of symptom days was reduced, in a healthy active population.

## Introduction

One of the leading causes of a visit to a general practitioner is upper respiratory tract infections (URTI), such as coughs, colds, pharyngitis and laryngitis [11]. It has also been demonstrated that every adult will suffer, on average, between 2 and 5 colds every year [17] with a large economic cost in lost work days and medical expenses. Several factors can alter the incidence of URTI with periods of high intensity/duration exercise training being shown to result in an approximately 2 fold increase in URTI incidence [21, 22]. This increase in URTI incidence has long been thought to be the result of perturbations in the function of the immune system, with many changes to immune function after both acute and chronic exercise [28]. What has remained challenging, however, has been establishing a clear link between individuals with the greatest perturbations of the immune system and URTI incidence. In this regard one of the most promising markers associated with URTI incidence is the levels of salivary immunoglobulin A (sIgA), with reduced sIgA having been demonstrated to be

associated with URTI incidence in athletic groups [10]. It is of clear importance, therefore, to uncover strategies for reducing URTI incidence in athletic populations with nutrition often being an area of focus.

In recent work, vitamin D has been shown to be crucial in the regulation of the immune system [1], with lower levels of vitamin D being associated with an increase in respiratory infections and decrease in lung function in a middle-aged British population [3]. Similar investigations have been carried out in athletic populations, with athletes with low vitamin D status having a higher risk of URTI and suffering more symptoms when a URTI bout is present [15]. There are as yet no vitamin D supplementation studies to investigate whether such a strategy may reduce URTI incidence in such a population. In a similar vein it is known that fish oil can improve immune function in the recovery period after exercise [13], suggesting that fish oil supplementation may reduce URTI incidence although, as with vitamin D, there are no studies investigating this at present. One method which has successfully been shown to reduce URTI incidence is through an

accepted after revision  
September 03, 2014

## Bibliography

DOI <http://dx.doi.org/10.1055/s-0034-1394464>  
Published online:  
February 9, 2015  
Int J Sports Med 2015; 36:  
426–430 © Georg Thieme  
Verlag KG Stuttgart · New York  
ISSN 0172-4622

## Correspondence

**Dr. Stuart Robert Gray, PhD**  
Institute of Medical Sciences  
University of Aberdeen  
Foresterhill  
Aberdeen  
United Kingdom AB25 2ZD  
Tel.: +44/122/4438 026  
Fax: +44/122/4555 719  
s.r.gray@abdn.ac.uk

increase in dietary protein intake [29]. In this study, with a counterbalanced design, participants performed 7 days of high-intensity exercise training on either a normal (1.5 g/kg body mass/day) or high-protein diet (3.0 g/kg body mass/day). The high protein diet attenuated the training-induced rise in URTI incidence and restored leukocyte trafficking. Although showing clear potential, this study was of short duration (7 days) and involved relatively small subject numbers (n=8) in a lab-based induction of an overreached state. Longer-term studies among larger athletic populations during their normal training regimen are therefore required.

The aim of the current study was to investigate the effects of supplementation with fish oil, vitamin D and protein on URTI incidence over a 16-week period in an athletic population and to determine whether such changes were associated with increases in sIgA.

## Methods



### Participants

42 participants (age 26.8±4.4 years; height 171.6±29.6 cm; body mass 68.2±14.8 kg and percentage body fat 15.4±6.9%; sex: 26 male, 16 female) were recruited to the study. 2 cohorts were recruited beginning in March 2013 (n=36) and September 2013 (n=6). All participants reported participating at least 3 h of moderate/high intensity exercise training per week. For the 16-week study period participants were asked to maintain their habitual exercise training and nutritional intake. The study was approved by the University of Aberdeen College of Life Sciences and Medicine Ethics Review Board, and participants were made aware of the aims, risk and discomfort associated with the study before providing written informed consent [14]. Participants were free to withdraw at any time.

### Supplementation

Participants were randomly assigned to either control (n=20; 9 females) or supplementation (n=22; 7 females) groups. Supplements consisted of drinks provided in unlabelled cartons, with participants being blinded as to which group they were assigned to. Following their initial visit, participants in the control group consumed 2 drinks per day, at a time of their choosing, containing water, sugar, lingonberry juice (6.2%), red grape juice (3.9%), ascorbic acid, citric acid and lingonberry aroma. Each drink was 200 ml with 100 kcal of energy (25 g carbohydrate, 0 g fat and 0 g protein). The supplement group consumed 2 drinks per day with the same ingredients, and each drink supplemented by 550 mg DHA, 550 mg EPA, 8 g whey protein isolate and 10 µg vitamin D3. Each drink was 200 ml with 200 kcal energy (28.5 g carbohydrate, 5.8 g fat and 8 g protein). Drinks were provided by Smartfish (Norway).

### Experimental protocol

Participants visited the laboratory in the morning at baseline and upon completion of the study. On both of these visits body mass was recorded to the nearest 0.1 kg using a weighing scale, and height was measured to the nearest 0.5 cm using a stadiometer. Skinfold thickness was measured with callipers at 4 sites (bicep, triceps, subscapular and suprailiac) to the nearest 0.1 mm on the right side of the body. Body fat percentage was calculated using standard equations [7]. During these visits, unstimulated saliva samples were also collected from 3 min of passive salivation.

Training load was monitored throughout the study period by asking participants to complete the standard short form of the International Physical Activity Questionnaire (IPAQ) every week. From this questionnaire, training load was calculated every week in metabolic equivalents (MET) min/week. Participants were also asked to complete a questionnaire to record URTI symptoms on a daily basis [8]. Participants were asked to take any medication, or visit their GP as normal when symptoms were present, with these events being recorded in their illness log. The illness log listed the following symptoms: sore throat, catarrh, runny nose, cough, fever, repetitive sneezing, joint aches and pains, general weakness, headache and loss of sleep. Severity ratings of mild, moderate and severe were given with the following guidance: mild=normal training, moderate=modified/reduced training and severe=discontinued training; and these were scored as 1, 2 and 3, respectively, providing a quantitative total symptom score for each day (calculated as symptoms multiplied by severity score). In any given week a symptom score of ≥12, and separated by at least one week from another week with a score ≥12, was taken as a single URTI episode. The total number of URTI symptom days was calculated as the number of days with a symptom score of ≥5 [23].

### Saliva IgA

The volume of saliva collected in 3 min was estimated by weighing, and the salivary flow rate was subsequently calculated. Saliva was analysed for IgA using a commercially available ELISA kit (Salimetrics, USA) according to the manufacturer's instructions. Samples and standards were analysed in duplicate with the intra-assay coefficient of variation being 7.9%. Secretion rate for IgA was calculated by multiplying the salivary flow rate by the saliva IgA concentration.

### Statistical analysis

Differences in the number of participants who suffered an URTI episode between control and supplemented groups were assessed by a chi-square test. To compare the number of URTI episodes and total number of symptom days, and, when a URTI was present, to compare URTI duration and symptom severity score, independent t-tests were performed. Changes in saliva IgA were compared using a two-way repeated measures ANOVA. Significance was accepted at p<0.05, and data are presented as mean±SD.

## Results




### Participants

42 participants were recruited to the study, with 6 participants fully withdrawing for a variety of reasons (e.g. injury, did not like the supplements, undisclosed reasons). A further 6 participants completed 12 weeks of the study, with 30 participants completing the full 16 weeks of the study and saliva samples and questionnaires being completed for all 36 of these participants. Body mass increased slightly in the control group during the intervention (68.2±14.5 kg at baseline to 69.2±14.4 kg upon completion) with a slight decrease in the supplemented group (71.1±9.8 kg at baseline to 69.2±14.4 kg upon completion). Similarly, percentage body fat increased slightly in the control group during the intervention (17.3±5.0% at baseline to 18.5±5.0% upon completion) with a slight decrease in the supplemented group (14.6±7.8% at baseline to 14.2±7.1% upon completion).

	Control (n = 17; 6 females)		Supplemented (n = 19; 6 females)	
	Baseline	16 weeks	Baseline	16 weeks
age (years)	28.3±6.4	–	26.7±4.4	–
height (cm)	174±11.9	–	177.9±6.9	–
body mass (kg)	68.2±14.5	69.2±14.4	71.1±9.8	70.7±6.9
body fat (%)	17.3±5.0	18.5±5.0	14.6±7.8	14.2±7.1
salivary IgA SR (µg/min)	64.5±36.8	69.1±30.2	63.2±31.0	73.1±32.7

SR = Secretion rate

**Table 1** Basic demographics and salivary IgA of participants at baseline and upon completion of the intervention.



**Fig. 1** Training load monitored via the IPAQ questionnaire on a weekly basis during the intervention.

These small changes were not significant with the repeated measures ANOVA, revealing no group, interaction or time effects in either body mass or percentage body fat (◊ **Table 1**).

### Training loads

The weekly training load, estimated via the IPAQ, was not different between the 2 groups with an average weekly load of  $3814 \pm 2108$  METmin/week in the control group and  $3893 \pm 2194$  METmin/week in the supplemented group (◊ **Fig. 1**).

### URTI incidence and severity

Data on URTI incidence and severity is presented in ◊ **Fig. 2**. There were no differences in the number of URTI episodes between groups. There were no differences in the percentage of participants reporting URTIs during the study period, with 49% of participants in the control group and 45% of participants in the supplemented group reporting a URTI episode. When a URTI was present there were no differences in either the severity or the duration of the episode. There were also no differences in the number of visits to a GP (1 participant in control and 2 in the supplemented group) or number of times medication was taken (medication taken 7 times in the control group and 5 in the supplemented group). However, the total number of symptom days reports was significantly lower in the supplement group compared to the control group ( $1.72 \pm 1.67$  in the supplemented group vs.  $2.79 \pm 1.76$  in the control group;  $<0.05$ ).

### Salivary IgA

◊ **Fig. 3** shows the sIgA concentration and secretion rate in control and supplemented groups upon entry to the study and upon completion of the intervention. There were no differences in


sIgA between visits or between the control and supplemented groups.

### Discussion


This 16-week study investigated the effects of a supplement containing fish oil, vitamin D and protein on self-reported URTI symptoms in a cohort of young active people. Throughout the trial, participants engaged in their normal training and competition schedule and were instructed to maintain their normal dietary habits. The main finding of the current study was that the supplement had no effect on the incidence and duration of URTI episodes, or the severity of symptoms when an URTI was present. Furthermore, sIgA concentration and secretion rate did not differ between groups. However, the supplemented group did report a lower number of symptom days compared to the control group. Taken together, this indicates that, while this supplement has no effect on URTI, the number of days with lower level symptoms, such as runny nose, cough, fever, catarrh, is reduced. As the guidance given to participants for scoring symptoms was related to the subsequent effect on training, this would still indicate that the supplement may be of use in maintaining training adherence in this population. Further work is clearly required to test this directly.

In the current study, these observed changes in symptom days were not associated with changes in sIgA, with no differences existing between groups either before or after the intervention. There is currently very little information on the effects of EPA, DHA, protein or vitamin D on sIgA levels, and, as the current study did not find any changes in URTI incidence, duration or severity, the lack of change in sIgA is perhaps not surprising. It has previously been shown that sIgA secretion rate does decrease from “optimal” to “inadequate” according to vitamin D status [15], although to our knowledge there have been no previous supplementation studies. With regard to protein supplementation, one study has demonstrated that glutamine supplementation is unable to attenuate the acute exercise-induced decrease in sIgA levels [18]. This, however, is the effect of a single amino acid on acute exercise responses. As such, the effect of chronic protein supplementation on sIgA levels remains to be tested. On the basis of the current study it appears unlikely that a very modest increase in protein would have any effect. Previous work has shown that in mice, fish oil supplementation can preserve bronchoalveolar IgA levels in response to infusion with 5-fluorouracil, an anticancer drug known to reduce mucosal immunity [9]. However, in a similar vein to protein it would appear likely that fish oil supplementation has very little effect on sIgA levels in humans.

The supplement investigated in the current study was chosen on the basis of the immunomodulatory effects of the individual components. While the current study is unable to determine



**Fig. 2** URTI data from participants in control and supplemented groups of the 16 week intervention period. **a** Average number of URTI episodes per participant, **b** Average duration of each URTI sustained, **c** Symptom severity score when URTI is sustained, **d** Total number of symptom days. \* denotes a significant difference between groups ( $P < 0.05$ ).



**Fig. 3** Salivary IgA (sIgA) concentration and secretion rate in control and supplemented groups at baseline and after 16 weeks of the intervention.

which of these supplements is responsible for the reduction in total symptom days, it is prudent to discuss the literature available on the effects of the individual supplements on the immune system.

Rich in EPA and DHA, fish oil has long been investigated for its effects on the immune system and its ability to reduce inflammation [4, 16, 25, 27]. Among exercise studies there are some conflicting results on the effects of fish oil consumption on such factors. Indeed, in a study by Toft and colleagues [26] it was found that fish oil supplementation did not alter plasma IL-6 levels after a marathon, with similar results being found after 3 days of intensive exercise [20]. However, we have recently demonstrated that fish oil supplementation can improve post-exercise peripheral blood mononuclear cell (PBMC) IL-2 production, NK cell activity and markers of oxidative stress [12, 13]. Thus far, no studies have investigated the effects of EPA and/or DHA on URTI incidence in active populations.

Historically known for its role in the absorption of calcium in the intestine, the physiological roles of vitamin D have expanded manifold in recent years [2]. For example, vitamin D has been demonstrated to promote antibacterial responses in monocytes/macrophages, altering T helper cell function and viral inactivation [24]. Consequently, the potential role of vitamin D in URTI incidence has been proposed due to the immune role of vitamin D. On this basis, He and colleagues [15] investigated the incidence of URTI compared to plasma vitamin D levels. This study demonstrated that participants who were deficient in vitamin D ( $< 30 \text{ nmol/L } 25(\text{O})\text{HD}$ ) had an increased number of symptom days and symptom severity score, with no significant effect on URTI incidence or duration. The findings do partly reflect the findings of the current study, indicating that supplementation with vitamin D may be contributing to the reduction in symptom days observed. However, a major drawback of the current study was that we did not measure plasma  $25(\text{O})\text{HD}$  and thus have no information as to whether our cohort were vitamin D deficient at baseline and it would seem reasonable to assume that any beneficial effects of vitamin D supplementation would be observed only in a deficient group.

With regard to the protein component of the supplement employed, it is well established that amino acids are crucial substrates for lymphocytes and an important modulator of their function [5]. Indeed it has been shown that increasing dietary

protein intake preserved neutrophil degranulation [6], superoxide production [19] and also that exercise-induced impairments in leukocyte trafficking are restored [29]. In this latter study the incidence of URTI was also reduced after increasing dietary protein intake during a period of high-intensity exercise training, suggesting that dietary protein may be of importance in determining URTI incidence. It is worth pointing out that in the study of Witard and colleagues, daily protein intake was increased by 118 g, which is far in excess of 16 g of protein that was given in the current study. It remains to be established whether this small increase in protein intake is a contributing factor to the reduction in symptom days observed in the current study.

As mentioned above, a major limitation of the current study was that baseline and post-interventional plasma levels of EPA, DHA and vitamin D were not measured. We are therefore unable to determine whether their levels were low at baseline or to monitor the increase post intervention.

In conclusion the current study has shown that a supplement containing fish oil, vitamin D and protein has no effect on the incidence or severity of URTI in a group of active people, although the total number of symptoms days was reduced. The current study was not designed to investigate the contributions of each of the components of this supplement, and further work is needed in this area and also in determining the optimal quantities of each supplement.

## Acknowledgements

This study was supported by Smartfish, Oslo, Norway.

**Conflict of interest:** The authors have no conflict of interest to declare.

## References

- Aranow C. Vitamin D and the immune system. *J Investig Med* 2011; 59: 881–886
- Basit S. Vitamin D in health and disease: A literature review. *Br J Biomed Sci* 2013; 70: 161–172
- Berry DJ, Hesketh K, Power C, Hypponen E. Vitamin D status has a linear association with seasonal infections and lung function in british adults. *Br J Nutr* 2011; 106: 1433–1440
- Calder PC, Yaqoob P, Thies F, Wallace FA, Miles EA. Fatty acids and lymphocyte functions. *Br J Nutr* 2002; 87 (Suppl 1): S31–S48
- Chuang JC, Yu CL, Wang SR. Modulation of lymphocyte proliferation by enzymes that degrade amino acids. *Clin Exp Immunol* 1990; 82: 469–472
- Costa RJS, Oliver SJ, Laing SJ, Walters R, Bilzon JJJ, Walsh NP. Influence of timing of postexercise carbohydrate-protein ingestion on selected immune indices. *Int J Sport Nutr Exerc Metab* 2009; 19: 366–384
- Durnin JV, Womersley J. Total body fat, calculated from body density, and its relationship to skinfold thickness in 571 people aged 12–72 years. *Proc Nutr Soc* 1973; 32: 45A
- Fricker PA, Pyne DB, Saunders PU, Cox AJ, Gleeson M, Telford RD. Influence of training loads on patterns of illness in elite distance runners. *Clin J Sport Med* 2005; 15: 246–252
- Fukatsu K, Nagayoshi H, Maeshima Y, Ueno C, Saitoh D, Mochizuki H. Fish oil infusion reverses 5-fluorouracil-induced impairments in mucosal immunity in mice. *Clinical Nutrition* 2008; 27: 269–275
- Gleeson M, Bishop N, Oliveira M, McCauley T, Tauler P, Muhamad AS. Respiratory infection risk in athletes: Association with antigen-stimulated IL-10 production and salivary IgA secretion. *Scand J Med Sci Sports* 2012; 22: 410–417
- Graham NH. The epidemiology of acute respiratory tract infections in children and adults: A global perspective. *Epidemiol Rev* 1990; 12: 149–178
- Gray P, Chappell A, Jenkinson AM, Thies F, Gray SR. Fish oil supplementation reduces markers of oxidative stress but not muscle soreness after eccentric exercise. *Int J Sport Nutr Exerc Metab* 2014; 24: 206–214
- Gray P, Gabriel B, Thies F, Gray SR. Fish oil supplementation augments post-exercise immune function in young males. *Brain, Behavior, and Immunity* 2012; 26: 1265–1272
- Harriss DJ, Atkinson G. Ethical standards in sport and exercise science research: 2014 update. *Int J Sports Med* 2013; 34: 1025–1028
- He CS, Handzlik M, Fraser WD, Muhamad AS, Preston H, Richardson A, Gleeson M. Influence of vitamin D status on respiratory infection incidence and immune function during 4 months of winter training in endurance sport athletes. *Exerc Immunol Rev* 2013; 19: 86–101
- Healy D, Wallace F, Miles E, Calder P, Newsholme P. Effect of low-to-moderate amounts of dietary fish oil on neutrophil lipid composition and function. *Lipids* 2000; 35: 763–768
- Heath GW, Macera CA, Nieman DC. Exercise and upper respiratory tract infections. is there a relationship? *Sports Med* 1992; 14: 353–365
- Krzywkowski K, Petersen EW, Ostrowski K, Link-Amster H, Boza J, Halkjaer-Kristensen J, Pedersen BK. Effect of glutamine and protein supplementation on exercise-induced decreases in salivary IgA. *Journal of Applied Physiology* 2001; 91: 832–838
- Nelson AR, Jackson L, Clarke J, Stellingwerff T, Broadbent S, Rowlands DS. Effect of post-exercise protein-leucine feeding on neutrophil function, immunomodulatory plasma metabolites and cortisol during a 6-day block of intense cycling. *Eur J Appl Physiol* 2013; 113: 2211–2222
- Nieman DC, Henson DA, McNulty SR, Jin F, Maxwell KR. N-3 polyunsaturated fatty acids do not alter immune and inflammation measures in endurance athletes. *Int J Sport Nutr Exerc Metab* 2009; 19: 536–546
- Nieman DC, Johanssen LM, Lee JW, Arabatzis K. Infectious episodes in runners before and after the los angeles marathon. *J Sports Med Phys Fitness* 1990; 30: 316–328
- Peters EM, Bateman ED. Ultramarathon running and upper respiratory tract infections. an epidemiological survey. *S Afr Med J* 1983; 64: 582–584
- Predy GN, Goel V, Lovlin R, Donner A, Stitt L, Basu TK. Efficacy of an extract of north american ginseng containing poly-furanosyl-pyranosyl-saccharides for preventing upper respiratory tract infections: A randomized controlled trial. *Canadian Medical Association Journal* 2005; 173: 1043–1048
- Priehl B, Treiber G, Pieber TR, Amrein K. Vitamin D and immune function. *Nutrients* 2013; 5: 2502–2521
- Rees D, Miles EA, Banerjee T, Wells SJ, Roynette CE, Wahle KW, Calder PC. Dose-related effects of eicosapentaenoic acid on innate immune function in healthy humans: A comparison of young and older men. *Am J Clin Nutr* 2006; 83: 331–342
- Toft AD, Thorn M, Ostrowski K, Asp S, Moller K, Iversen S, Hermann C, Sondergaard SR, Pedersen BK. N-3 polyunsaturated fatty acids do not affect cytokine response to strenuous exercise. *J Appl Physiol* 2000; 89: 2401–2406
- Tull SP, Yates CM, Maskrey BH, O'Donnell VB, Madden J, Grimble RF, Calder PC, Nash GB, Rainger GE. Omega-3 fatty acids and inflammation: Novel interactions reveal a new step in neutrophil recruitment. *PLoS Biol* 2009; 7: e1000177
- Walsh NP, Gleeson M, Shephard RJ, Gleeson M, Woods JA, Bishop NC, Fleschner M, Green C, Pedersen BK, Hoffman-Goetz L, Rogers CJ, Northoff H, Abbasi A, Simon P. Position statement. part one: Immune function and exercise. *Exerc Immunol Rev* 2011; 17: 6–63
- Witard OC, Turner JE, Jackman SR, Kies AK, Jeukendrup AE, Bosch JA, Tipton KD. High dietary protein restores overreaching induced impairments in leukocyte trafficking and reduces the incidence of upper respiratory tract infection in elite cyclists. *Brain Behav Immun* 2014; 39: 211–219